Condition	Category	Measure Name
Surgery patients who were taking heart drugs called beta blockers before coming to the hospital, who were kept on the beta blockers during the period just before and after their surgery	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Surgery patients who were given the right kind of antibiotic to help prevent infection	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Heart surgery patients whose blood sugar (blood glucose) is kept under good control in the days right after surgery	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Surgery patients needing hair	Surgical Care Improvement	National Average of Hospitals


Page 1 of 22 05/30/2011



Page 2 of 22 05/30/2011

removed from the surgical area before surgery, who had hair removed using a safer method (electric clippers or hair removal cream – not a razor)	Project	submitting data:
Surgery patients whose doctors ordered treatments to prevent blood clots after certain types of surgeries	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Patients who got treatment at the right time (within 24 hours before or after their surgery) to help prevent blood clots after certain types of surgery	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Heart Attack Patients Given Aspirin at Arrival	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given Aspirin at Discharge	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given Smoking Cessation Advice/Counseling	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given	Heart Attack or Chest Pain	National Average of Hospitals


Page 3 of 22 05/30/2011



Page 4 of 22 05/30/2011

Beta Blocker at Discharge		submitting data:
Heart Attack Patients Given Fibrinolytic Medication Within 30 Minutes Of Arrival	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Heart Attack Patients Given PCI Within 90 Minutes Of Arrival	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Pneumonia Patients Assessed and Given Pneumococcal Vaccination	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Given Smoking Cessation Advice/Counseling	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Given Initial Antibiotic(s) within 6 Hours After Arrival	Pneumonia	National Average of Hospitals submitting data:
Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)	Pneumonia	National Average of Hospitals submitting data:


Page 5 of 22 05/30/2011



Page 6 of 22 05/30/2011

Pneumonia Patients Assessed and Given Influenza Vaccination	Pneumonia	National Average of Hospitals submitting data:
Heart Failure Patients Given Discharge Instructions	Heart Failure	National Average of Hospitals submitting data:
Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function	Heart Failure	National Average of Hospitals submitting data:
Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Failure	National Average of Hospitals submitting data:
Heart Failure Patients Given Smoking Cessation Advice/Counseling	Heart Failure	National Average of Hospitals submitting data:
Children Who Received Reliever Medication While Hospitalized for Asthma	Children's Asthma	National Average of Hospitals submitting data:
Children Who Received Systemic Corticosteroid Medication (oral and IV Medication That Reduces Inflammation and Controls Symptoms) While Hospitalized for Asthma	Children's Asthma	National Average of Hospitals submitting data:
Children and their Caregivers	Children's Asthma	National Average of Hospitals


Page 7 of 22 05/30/2011



Page 8 of 22 05/30/2011

Who Received a Home Management Plan of Care Document While Hospitalized for Asthma		submitting data:
Outpatients having surgery who got an antibiotic at the right time - within one hour before surgery (higher numbers are better)	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Outpatients having surgery who got the right kind of antibiotic (higher numbers are better)	Surgical Care Improvement Project	National Average of Hospitals submitting data:
Average number of minutes before outpatients with chest pain or possible heart attack who needed specialized care were transferred to another hospital (a lower number of minutes is better)	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Average number of minutes before outpatients with chest	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
pain or possible heart attack got an ECG (a lower number of minutes is better)		

Page 9 of 22 05/30/2011



Page 10 of 22 05/30/2011

possible heart attack who got drugs to break up blood clots within 30 minutes of arrival (higher numbers are better)		submitting data:
Outpatients with chest pain or possible heart attack who got aspirin within 24 hours of arrival (higher numbers are better)	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Median Time to Fibrinolysis	Heart Attack or Chest Pain	National Average of Hospitals submitting data:
Surgery patients who were taking heart drugs called beta blockers before coming to the hospital, who were kept on the beta blockers during the period just before and after their surgery	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients who were given the right kind of antibiotic to help prevent infection	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than

Page 11 of 22 05/30/2011



Page 12 of 22 05/30/2011

Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Heart surgery patients whose blood sugar (blood glucose) is kept under good control in the days right after surgery	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients needing hair removed from the surgical area before surgery, who had hair removed using a safer method (electric clippers or hair removal cream – not a razor)	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Surgery patients whose doctors ordered treatments to prevent blood clots after certain types of surgeries	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Patients who got treatment at the right time (within 24 hours before or after their surgery) to help prevent blood clots after certain types of surgery	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Aspirin at Arrival	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data


Page 13 of 22 05/30/2011



Page 14 of 22 05/30/2011

Aspirin at Discharge		scored equal to or higher than
Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Smoking Cessation Advice/Counseling	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Beta Blocker at Discharge	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given Fibrinolytic Medication Within 30 Minutes Of Arrival	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Attack Patients Given PCI Within 90 Minutes Of Arrival	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Assessed and Given Pneumococcal Vaccination	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than


Page 15 of 22 05/30/2011



Page 16 of 22 05/30/2011

Pneumonia Patients Given Smoking Cessation Advice/Counseling	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Given Initial Antibiotic(s) within 6 Hours After Arrival	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Pneumonia Patients Assessed and Given Influenza Vaccination	Pneumonia	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given Discharge Instructions	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than
Heart Failure Patients Given Smoking Cessation Advice/Counseling	Heart Failure	Top 10% of Hospitals submitting data scored equal to or higher than
Outpatients having surgery who got an antibiotic at the	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than


Page 17 of 22 05/30/2011



Page 18 of 22 05/30/2011

who got an antibiotic at the right time - within one hour before surgery (higher numbers are better)	Project	scored equal to or higher than
Outpatients having surgery who got the right kind of antibiotic (higher numbers are better)	Surgical Care Improvement Project	Top 10% of Hospitals submitting data scored equal to or higher than
Average number of minutes before outpatients with chest pain or possible heart attack who needed specialized care were transferred to another hospital (a lower number of minutes is better)	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Average number of minutes before outpatients with chest pain or possible heart attack got an ECG (a lower number of minutes is better)	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Outpatients with chest pain or possible heart attack who got drugs to break up blood clots within 30 minutes of arrival (higher numbers are better)	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data scored equal to or higher than
Outpatients with chest pain or	Heart Attack or Chest Pain	Top 10% of Hospitals submitting data

Page 19 of 22 05/30/2011



Page 20 of 22 05/30/2011

possible heart attack who got aspirin within 24 hours of arrival (higher numbers are better) scored equal to or higher than

Median Time to Fibrinolysis

Heart Attack or Chest Pain

Top 10% of Hospitals submitting data scored equal to or higher than

Page 21 of 22 05/30/2011

20

Page 22 of 22 05/30/2011